

or a stereoisomeric form of the compound of formula I, or a physiologically tolerated salt of the compound of formula I; and

a third component comprising a pharmaceutically tolerated excipient;

wherein the first component has a concentration from about 2 to about 20 mg and the second component has a concentration from about 0.3% to about 50% of the first component.



20. A method of treating an immunological disease comprising administering to a patient in need of such treatment, a therapeutically effective amount of a solid composition comprising

a first component comprising 5-methyl-4'-trifluoromethyl-4-isoxazolecarboxanilide;

a second component comprising a compound of formula I

LAW OFFICES

FINNEGAN, HENDERSON,  
FARABOW, GARRETT,  
& DUNNER, L.L.P.  
1300 I STREET, N.W.  
WASHINGTON, DC 20005  
202-408-4000



or a stereoisomeric form of the compound of formula I, or a physiologically tolerated salt of the compound of formula I; and

a third component comprising a pharmaceutically tolerated excipient;

wherein the first component has a concentration from about 2 to about 20 mg and the second component has a concentration from about 0.3% to about 50% of the first component.

26. A method of treating a disease comprising administering to a patient in need of such treatment, a therapeutically effective amount of a solid composition comprising

a first component comprising 5-methyl-4'-trifluoromethyl-4-isoxazolecarboxanilide;



a second component comprising a compound of formula I

05101572.102398

LAW OFFICES

FINNEGAN, HENDERSON,  
FARABOW, GARRETT  
& DUNNER, L. L. P.  
1300 I STREET, N. W.  
WASHINGTON, D. C. 20005  
202-408-4000

B3



or a stereoisomeric form of the compound of formula I, or a physiologically tolerated salt of the compound of formula I; and

a third component comprising a pharmaceutically tolerated excipient;

wherein the first component has a concentration from about 2 to about 20 mg and the second component has a concentration from about 0.3% to about 50% of the first component, and wherein the disease is atopic dermatitis, asthma, urticaria, rhinitis, uveitis, type II diabetes, cystic fibrosis, colitis, or hepatic fibrosis.

27. A method of treating a cancerous disease comprising administering to a therapeutically effective amount of a solid composition comprising
- a first component comprising 5-methyl-4'-trifluoromethyl-4-isoxazolecarboxanilide;
  - a second component comprising a compound of formula I